

# Numbers of Americans Affected by Mental Illness

- One in four adults-approximately 61.5 million Americans-experience mental illness in a given year. One in 17-about 13.6 million-live with a serious mental illness such as schizophrenia, major depression or bipolar disorder.<sup>1</sup>
- Approximately 20 percent of youth ages 13 to 18 experience severe mental disorders in a given year. For ages 8 to 15, the estimate is 13 percent.<sup>2</sup>
- Approximately 1.1 percent of American adults about 2.6 million people—live with schizophrenia.<sup>3,4</sup>
- Approximately 2.6 percent of American adults–6.1 million people–live with bipolar disorder.
- Approximately 6.7 percent of American adults–about 14.8 million people–live with major depression.<sup>4,6</sup>
- Approximately 18.1 percent of American adults–about 42 million people–live with anxiety disorders, such as panic disorder, obsessive-compulsive disorder (OCD), posttraumatic stress disorder (PTSD), generalized anxiety disorder and phobias.<sup>4,7</sup>
- About 9.2 million adults have co-occurring mental health and addiction disorders.<sup>8</sup>
- Approximately 26 percent of homeless adults staying in shelters live with serious mental illness and an estimated 46 percent live with severe mental illness and/or substance use disorders.<sup>9</sup>
- Approximately 20 percent of state prisoners and 21 percent of local jail prisoners have "a recent history" of a mental health condition.<sup>10</sup>
- Seventy percent of youth in juvenile justice systems have at least one mental health condition and at least 20 percent live with a severe mental illness.<sup>11</sup>

### **Getting Mental Health Treatment in America**

• Approximately 60 percent of adults<sup>12</sup>, and almost one-half

## Mental Illness FACTS AND NUMBERS

of youth ages 8 to 15 with a mental illness received no mental health services in the previous year.<sup>13</sup>

- African American and Hispanic Americans used mental health services at about one-half the rate of whites in the past year and Asian Americans at about one-third the rate.<sup>14.</sup>
- One-half of all chronic mental illness begins by the age of 14; three-quarters by age 24.<sup>15</sup> Despite effective treatment, there are long delays—sometimes decades—between the first appearance of symptoms and when people get help.<sup>16</sup>

### The Impact of Mental Illness in America

- Serious mental illness costs America \$193.2 billion in lost earnings per year.<sup>17</sup>
- Mood disorders such as depression are the third most common cause of hospitalization in the U.S. for both youth and adults ages 18 to 44.<sup>18</sup>
- Individuals living with serious mental illness face an increased risk of having chronic medical conditions.
  Adults living with serious mental illness die on average 25 years earlier than other Americans, largely due to treatable medical conditions.
- Over 50 percent of students with a mental health condition age 14 and older who are served by special education drop out-the highest dropout rate of any disability group.<sup>21</sup>
- Suicide is the tenth leading cause of death in the U.S. (more common than homicide) and the third leading cause of death for ages 15 to 24 years.<sup>22</sup> More than 90 percent of those who die by suicide had one or more mental disorders.<sup>23</sup>
- Although military members comprise less than 1 percent of the U.S. population<sup>24</sup>, veterans represent 20 percent of suicides nationally. Each day, about 18 veterans die from suicide.<sup>25</sup>

NAMI • The National Alliance on Mental Illness • 1 (800) 950-NAMI • www.nami.org 3803 N. Fairfax Drive, Suite 100, Arlington, VA 22203



#### References

1 National Institutes of Health, National Institute of Mental Health. (n.d.). *Statistics: Any Disorder Among Adults*. Retrieved March 5, 2013, from <a href="http://www.nimh.nih.gov/statistics/1ANYDIS\_ADULT.shtml">http://www.nimh.nih.gov/statistics/1ANYDIS\_ADULT.shtml</a>

2 National Institutes of Health, National Institute of Mental Health. (n.d.). *Any Disorder Among Children*. Retrieved March 5, 2013, from <u>http://www.nimh.nih.gov/statistics/1ANYDIS\_CHILD.shtml</u>

3 National Institutes of Health, National Institute of Mental Health. (n.d.) *The Numbers Count: Mental Disorders in America*. Retrieved March 5, 2013, from <a href="http://www.nimh.nih.gov/health/publications/the-numbers-count-mental-disorders-in-america/index.shtml">http://www.nimh.nih.gov/health/publications/the-numbers-count-mental-disorders-in-america/index.shtml</a>

4 Prevalence numbers were calculated using NIMH percentages (cited) and 2010 Census data. Census data is available at: United States Census Bureau. (revised 2011). "USA [State & County QuickFacts]." Retrieved March 5, 2013, from <u>http://quickfacts.census.gov/qfd/states/00000.html</u> 5 National Institutes of Health, National Institute of Mental Health. (n.d.). *The Numbers Count: Mental Disorders in America*. Retrieved March 5, 2013, from <u>http://www.nimh.nih.gov/health/publications/the-numbers-count-mental-disorders-in-america/index.shtml</u> 6 *Ibid*.

7 National Institutes of Health, National Institute of Mental Health. (n.d.). *Statistics: Any Anxiety Disorder Among Adults*. Retrieved March 5, 2013, from <a href="http://www.nimh.nih.gov/statistics/lanyanx\_adult.shtml">http://www.nimh.nih.gov/statistics/lanyanx\_adult.shtml</a>

8 Substance Abuse and Mental Health Services Administration. (2012). *Results from the 2010 National Survey on Drug Use and Health: Mental Health Findings* NSDUH Series H-42, HHS Publication No. (SMA) 11-4667). Rockville, Md.: Substance Abuse and Mental Health Services Administration, 2012.

9 U.S. Department of Housing and Urban Development, Office of Community Planning and Development. (2011). *The 2010 Annual Homeless Assessment Report to Congress*. Retrieved March 5, 2013, from <u>http://www.hudhre.info/documents/2010HomelessAssessmentReport.pdf</u> 10 Glaze, L.E. & James, D.J. (2006, updated December). Mental Health Problems of Prison and Jail Inmates. *Bureau of Justice Statistics Special Report*. U.S. Department of Justice, Office of Justice Programs Washington, D.C. Retrieved March 5, 2013, from

http://bjs.ojp.usdoj.gov/content/pub/pdf/mhppji.pdf

11 Skowyra, K.R. & Cocozza, J.J. (2007) Blueprint for Change: A Comprehensive Model for the Identification and Treatment of Youth with Mental Health Needs in Contact with the Juvenile Justice System. The National Center for Mental Health and Juvenile Justice; Policy Research Associates, Inc. The Office of Juvenile Justice and Delinquency Prevention. Delmar, N.Y: The National Center for Mental Health and Juvenile Justice; Policy Research Associates, Inc.

12Substance Abuse and Mental Health Services Administration. (2012). *Results from the 2010 National Survey on Drug Use and Health: Mental Health Findings* NSDUH Series H-42, HHS Publication No. (SMA) 11-4667). Rockville, Md.; Substance Abuse and Mental Health Services Administration, 2012.

13 National Institute of Mental Health. (n.d.). Use of Mental Health Services and Treatment Among Children. Retrieved March 5, 2013, from <a href="http://www.nimh.nih.gov/statistics/1NHANES.shtml">http://www.nimh.nih.gov/statistics/1NHANES.shtml</a>

14 Ibid.

15 Kessler, R.C, et al. (2005). Lifetime prevalence and age-of-onset distributions of *DSM-IV* disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62(6), 593-602.

16 National Institutes of Health, National Institute of Mental Health. (2005). *Mental Illness Exacts Heavy Toll, Beginning in Youth*. Retrieved March 5, 2013, from <a href="http://www.nih.gov/news/pr/jun2005/nimh-06.htm">http://www.nih.gov/news/pr/jun2005/nimh-06.htm</a>

17 Insel, T.R. (2008). Assessing the Economic Costs of Serious Mental Illness. The American Journal of Psychiatry. 165(6), 663-665.

18 Wier, LM (Thompson Reuters), et al. *HCUP facts and figures: statistics on hospital-based care in the United States, 2009.* Web.. Rockville, Md. Agency for Healthcare Research and Quality, 2011. Retrieved March 5, 2013, from http://www.hcup-us.ahrq.gov/reports.jsp.

19 Colton, C.W. & Manderscheid, R.W. (2006). Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Preventing Chronic Disease: Public Health Research, Practice and Policy*, 3(2), 1-14.

20 Parks, J., et al. (2006). *Morbidity and Mortality in People with Serious Mental Illness*. Alexandria, VA: National Association of State Mental Health Program Directors (NASMHPD) Medical Directors Council.

21 U.S. Department of Education. (2006). Twenty-eighth annual report to Congress on the implementation of the Individuals with Disabilities Education Act, 2006, Vol. 2. Washington, D.C.: U.S. Department of Education.

22 McIntosh, J.L.. & Drapeau, C.W. (for the American Association of Suicidology). (2012). U.S.A. suicide: 2010

official final data. Washington, D.C: American Association of Suicidology.

23 American Association of Suicidology. (2012). *Suicide in the USA Based on 2010 Data*. Washington, DC: American Association of Suicidology. 24 Martinez, L. & Bingham, A. (2011). U.S. Veterans: by the Numbers. Retrieved March 5, 2013, from <u>http://abcnews.go.com/Politics/us-veterans-numbers/story?id=14928136</u>

25 Department of Veterans Affairs, Office of Patient Care Services, Office of Mental Health Services. (2011). *Fact Sheet VA Suicide Prevention Program Facts about Veteran Suicide*. Retrieved March 5, 2013, from <a href="https://www.erie.va.gov/pressreleases/assets/SuicidePreventionFactSheet.doc">www.erie.va.gov/pressreleases/assets/SuicidePreventionFactSheet.doc</a>

Reviewed by Ken Duckworth, M.D., March 2013

NAMI • The National Alliance on Mental Illness • 1 (800) 950-NAMI • www.nami.org 3803 N. Fairfax Drive, Suite 100, Arlington, VA 22203